arXiv:hep-th/9902081v1 11 Feb 1999

DAMTP 98-157
APCTP-1999004
hep-th/9902081

A Study of a Non-Abelian Generalization of the Born-Infeld Action

Jeong-Hyuck Park*

Asia Pacific Center for Theoretical Physics
207-43 Cheongryangri-dong Dongdaemun-gu
Seoul 130-012, Korea

Abstract

A new type of non-Abelian generalization of the Born-Infeld action is proposed, in which the spacetime indices and group indices are combined. The action is manifestly Lorentz and gauge invariant. In its power expansion, the lowest order term is the Yang-Mills action and the second term corresponds to the bosonic stringy correction to this action. Solutions of the Euler-Lagrange equation for the SU(2) case are considered and we show that there exists an instanton-like solution which has winding number one and finite action.

---

*E-mail address: J.H.Park@damtp.cam.ac.uk

1 Introduction

The Born-Infeld action [1] has been of renewed attention for reasons connected with M/string theory, but perhaps interest has also been enhanced by its compact and elegant form. There have been several proposals for the non-Abelian generalization of the Born-Infeld action by taking either a trace [2], an anti-symmetrized trace [3] or a symmetrized trace [4] of the group indices after formally expanding the ordinary Born-Infeld Lagrangian with the non-Abelian gauge fields substituted. In particular, the anti-symmetrized trace structure generates only odd powers of the field strength, F, while the symmetrized trace structure gives only even powers. Bosonic stringy corrections to the Yang-Mills action give a F^3 term at the lowest order [5] while for superstring theory the cubic term is absent [6,7]. However, the issue as to what is the natural non-Abelian generalization of the Born-Infeld action does not seem to have been resolved fully yet [8].

In this paper, we propose a different non-Abelian generalization of the Born-Infeld action for SU(N) gauge fields, where the spacetime indices and group indices are combined together. The action is manifestly Lorentz and gauge invariant. In its power expansion, the lowest order term is the Yang-Mills action and the second term corresponds to the bosonic stringy correction to this action. Solutions of the Euler-Lagrange equation for the SU(2) case are considered[1] and we show that there exists an instanton-like solution which has winding number one and finite action.

2 Non-Abelian Born-Infeld Lagrangian

The generators of the SU(N) group in the fundamental representation, $T^a = -T^{a\dagger}$, $\operatorname{tr} T^a = 0$, $a = 1, 2, \cdots, N^2 - 1$, may be chosen conventionally to satisfy

$$T^a T^b = -\tfrac{1}{2N}\delta^{ab} + \tfrac{1}{2}(id^{abc} + f^{abc})T^c \tag{2.1}$$

where d^{abc}, f^{abc} are real and totally symmetric, anti-symmetric over the indices, a, b, c. Eq.(2.1) implies

$$\operatorname{tr}(T^a T^b) = -\tfrac{1}{2}\delta^{ab} \tag{2.2}$$

and a completeness relation follows from the fact that $\{1, T^a\}$ forms a basis of $N \times N$ matrices

$$\sum_a T^a_{\alpha\beta} T^a_{\gamma\delta} = \tfrac{1}{2N}\delta_{\alpha\beta}\delta_{\gamma\delta} - \tfrac{1}{2}\delta_{\beta\gamma}\delta_{\alpha\delta} \tag{2.3}$$

---

[1] While this work was being completed, monopole solutions for the symmetrized non-Abelian Born-Infeld action were discussed in [9].

The gauge field, A_μ, is Lie algebra valued

$$A_\mu = A^a_\mu T^a \tag{2.4}$$

and the field strength is given by

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + [A_\mu, A_\nu] \tag{2.5}$$

In this paper, we regard the field strength, F, as a $Nd \times Nd$ matrix acting on $\mathbb{R}^N_{\text{gauge}} \otimes \mathbb{R}^d_{\text{spacetime}}$, where d is the spacetime dimension, by pairing the spacetime index, μ, and group index, α, together

$$F_{\mu\alpha,\,\nu\beta} \equiv F^a_{\mu\nu} T^a_{\alpha\beta} \tag{2.6}$$

We also expand the spacetime metric, $g_{\mu\nu}$, to a $Nd \times Nd$ matrix

$$g_{\mu\nu} \longrightarrow G_{\mu\alpha,\,\nu\beta} \equiv g_{\mu\nu}\delta_{\alpha\beta} \tag{2.7}$$

The non-Abelian Yang-Mills kinetic term is then

$$-\tfrac{1}{4} F^a_{\mu\nu} F^{a\mu\nu} = -\tfrac{1}{2}\mathrm{Tr}\mathcal{F}^2 \tag{2.8}$$

where $\mathcal{F}$ is a $Nd \times Nd$ matrix given by

$$\mathcal{F} \equiv G^{-1} F \tag{2.9}$$

Tr and Det indicate the trace and determinant of $Nd \times Nd$ matrices.
The proposed non-Abelian generalization of the Born-Infeld Lagrangian is then

$$\begin{aligned}
\mathcal{L} &= 2N\kappa^{-2}\left((\mathrm{Det}(G+\kappa F))^{\frac{1}{2N}} - \sqrt{|g|}\right) \\
&= 2N\kappa^{-2}\sqrt{|g|}\left(\exp\left(\textstyle\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n}\frac{1}{2N}\kappa^n \mathrm{Tr}\mathcal{F}^n\right) - 1\right) \\
&= \sqrt{|g|}\left(-\tfrac{1}{2}\mathrm{Tr}\mathcal{F}^2 + \tfrac{1}{3}\kappa\mathrm{Tr}\mathcal{F}^3 - \tfrac{1}{4}\kappa^2\mathrm{Tr}\mathcal{F}^4 + \tfrac{1}{16N}\kappa^2(\mathrm{Tr}\mathcal{F})^2 + \cdots\right)
\end{aligned} \tag{2.10}$$

where κ is a coupling constant. In the second line we have used the identity, $\mathrm{Det}M = \exp(\mathrm{Tr}\ln M)$.
The Lagrangian, $\mathcal{L}$, (2.10) is clearly real and gauge invariant.[2] The lowest order term or

---

[2] Note that G, F are hermitian.

$\kappa \to 0$ limit corresponds to the ordinary Yang-Mills action (2.8).[3]
From eq.(2.1) we can calculate the traces appearing in eq.(2.10)

$$\begin{aligned}\mathcal{L} = \;& \sqrt{|g|}\Big[-\tfrac{1}{4}F^a_{\mu\nu}F^{a\mu\nu} - \tfrac{1}{12}\kappa f^{abc}F^{a\mu}{}_{\nu}F^{b\nu}{}_{\lambda}F^{c\lambda}{}_{\mu} \\ & +\tfrac{1}{64N}\kappa^2\Big((F^a_{\mu\nu}F^{a\mu\nu})^2 - (4\delta^{ab}\delta^{cd} + 2Nd^{abe}d^{cde} - 2Nf^{abe}f^{cde})F^{a\mu}{}_{\nu}F^{b\nu}{}_{\lambda}F^{c\lambda}{}_{\rho}F^{d\rho}{}_{\mu}\Big) \\ & +\cdots\Big]\end{aligned} \tag{2.11}$$

From now on we restrict to flat spacetime with the metric η.
If we define[4]

$$\begin{aligned}\Pi^{a\mu\nu} &= -\Pi^{a\nu\mu} \\ &= -2\frac{\partial\mathcal{L}(F)}{\partial F^a_{\mu\nu}} = \kappa^{-1}(\mathrm{Det}(G+\kappa F))^{\frac{1}{2N}}\Big((G+\kappa F)^{-1\mu\alpha,\,\nu\beta} - (G+\kappa F)^{-1\nu\alpha,\,\mu\beta}\Big)T^a_{\beta\alpha} \\ &= F^{a\mu\nu} - \tfrac{1}{2}\kappa f^{abc}F^{b\mu\lambda}F^{c}{}_{\lambda}{}^{\nu} + \cdots\end{aligned} \tag{2.12}$$

then the equation of motion is

$$0 = D_\mu\Pi^{\mu\nu} = \partial_\mu\Pi^{\mu\nu} + [A_\mu, \Pi^{\mu\nu}] \qquad\qquad \Pi^{\mu\nu} = \Pi^{a\mu\nu}T^a \tag{2.13}$$

The energy-momentum tensor may be given either by Noether procedure

$$\begin{aligned}T^{\mu\nu}_N &= \frac{\partial\mathcal{L}}{\partial A^a_{\lambda,\mu}}A^{a,\nu}_{\lambda} - \eta^{\mu\nu}\mathcal{L} \\ &= -\Pi^{a\mu\lambda}A^{a,\nu}_{\lambda} - \eta^{\mu\nu}\mathcal{L}\end{aligned} \tag{2.14}$$

---

[3] We also note that the cubic term is identical with the bosonic stringy correction to the Yang-Mills action [5].

[4] We define $\frac{\partial}{\partial F^a_{\mu\nu}}J^{b\lambda\rho}F^b_{\lambda\rho} = \frac{1}{2}(J^{a\mu\nu} - J^{a\nu\mu})$.

or by taking variation with respect to the metric

$$\begin{aligned} T_g^{\mu\nu} &= -\frac{2}{\sqrt{|g|}}\frac{\partial \mathcal{L}}{\partial g_{\mu\nu}}|_{g=\eta} \\ &= -\kappa^{-2}\left(\mathrm{Det}(G+\kappa F)\right)^{\frac{1}{2N}}\left((G+\kappa F)^{-1\mu\alpha,\,\nu\alpha}+(G+\kappa F)^{-1\nu\alpha,\,\mu\alpha}\right)|_{g=\eta}+2N\kappa^{-2}\eta^{\mu\nu} \\ &= \Pi^{a\mu\lambda}F_{\lambda}^{a\nu}-\eta^{\mu\nu}\mathcal{L} \end{aligned} \tag{2.15}$$

The last equality in eq.(2.15) comes from observing

$$(G+\kappa F)^{-1}=G^{-1}-\kappa(G+\kappa F)^{-1}FG^{-1}=G^{-1}-\kappa G^{-1}F(G+\kappa F)^{-1} \tag{2.16}$$

The difference between $T_N^{\mu\nu}$ and $T_g^{\mu\nu}$ is with eq.(2.13)

$$T_N^{\mu\nu}-T_g^{\mu\nu}=\partial_\lambda(\Pi^{a\lambda\mu}A^{a\nu}) \tag{2.17}$$

which is consistent with the energy-momentum tensor conservation

$$\partial_\mu T_N^{\mu\nu}=\partial_\mu T_g^{\mu\nu}=0 \tag{2.18}$$

3 SU(2) Action in Four-dimensional Euclidean Space

We now consider the gauge group SU(2) and also restrict to four-dimensional Euclidean space, $x^0 \rightarrow -ix^4$, $\eta_{\mu\nu}=\delta_{\mu\nu}$ so that upper and lower Lorentz indices may be identified. We take $T^a=-\frac{i}{2}\sigma^a$, $f^{abc}=\epsilon^{abc}$, where σ^a, $a=1,2,3$ are Pauli matrices.
If we define $E_{\mu\alpha,\,\nu\beta}=\delta_{\mu\nu}\epsilon_{\alpha\beta}$ with a 2×2 anti-symmetric matrix, ϵ, then from $\epsilon\sigma^a\epsilon=(\sigma^a)^t$, $E(G+\kappa F)$ is anti-symmetric so that

$$\mathrm{Det}(G+\kappa F)=(\mathrm{Pfaffian}(EG+\kappa EF))^2 \tag{3.1}$$

Using this for the SU(2) case, the Lagrangian (2.10), $\mathcal{L}\rightarrow -\mathcal{L}_{\mathrm{E}}$, becomes

$$\begin{aligned} \mathcal{L}_{\mathrm{E}} &= 4\kappa^{-2}\left[1-\left(1-\tfrac{1}{4}\kappa^2\mathrm{Tr}\mathcal{F}^2+\tfrac{1}{6}\kappa^3\mathrm{Tr}\mathcal{F}^3-\tfrac{1}{8}\kappa^4\mathrm{Tr}\mathcal{F}^4+\tfrac{1}{32}\kappa^4(\mathrm{Tr}\mathcal{F}^2)^2\right)^{\frac{1}{2}}\right] \\ &= 4\kappa^{-2}\left[1-\left(1-\tfrac{1}{8}\kappa^2F^a_{\mu\nu}F^a_{\mu\nu}-\tfrac{1}{24}\kappa^3\epsilon^{abc}F^a_{\mu\nu}F^b_{\nu\lambda}F^c_{\lambda\mu}+\tfrac{1}{128}\kappa^4(F^a_{\mu\nu}F^a_{\mu\nu})^2\right.\right. \\ &\qquad \left.\left.-\tfrac{1}{32}\kappa^4F^a_{\mu\nu}F^a_{\nu\lambda}F^b_{\lambda\rho}F^b_{\rho\mu}+\tfrac{1}{64}\kappa^4F^a_{\mu\nu}F^b_{\nu\lambda}F^a_{\lambda\rho}F^b_{\rho\mu}\right)^{\frac{1}{2}}\right] \end{aligned} \tag{3.2}$$

Now we seek an SO(4) invariant solution in four-dimensional Euclidean space by adopting the ansatz of [10]

$$A_\mu(x) = f(r) g(x)^{-1} \partial_\mu g(x) \qquad g(x) = \frac{1}{r}(x_4 - i x_i \sigma^i) \in \mathrm{SU}(2) \tag{3.3}$$

where $r = \sqrt{\mathrm{x}^2 + x_4^2}$.
After some calculation, we get

$$\begin{aligned} F_{\mu\nu} &= \left(X(\delta_{\mu\lambda}\delta_{\nu\rho} - \delta_{\mu\rho}\delta_{\nu\lambda}) + Y \epsilon_{\mu\nu\lambda\rho}\right) x_\lambda g^{-1} \partial_\rho g \\ {}^*F_{\mu\nu} &= \tfrac{1}{2}\epsilon_{\mu\nu\lambda\rho} F_{\lambda\rho} = \left(Y(\delta_{\mu\lambda}\delta_{\nu\rho} - \delta_{\mu\rho}\delta_{\nu\lambda}) + X \epsilon_{\mu\nu\lambda\rho}\right) x_\lambda g^{-1} \partial_\rho g \end{aligned} \tag{3.4}$$

and

$$\Pi_{\mu\nu} = (1 - 2\kappa Y - 3\kappa^2 X^2)^{-\frac{1}{2}} (F_{\mu\nu} + \kappa X {}^*F_{\mu\nu}) \tag{3.5}$$

where

$$X = \frac{f'}{r} \qquad Y = \frac{2}{r^2}(f-1)f \tag{3.6}$$

Using the Bianchi identity

$$D_\mu {}^*F_{\mu\nu} = 0 \tag{3.7}$$

and

$$D_\mu F_{\mu\nu} = (f'' + X + 2Y - 4fY) g^{-1} \partial_\nu g \tag{3.8}$$

the equation of motion (2.13) gives a second order differential equation

$$0 = f''(1+\kappa Y)(1-2\kappa Y) + 2(2f-1)(\kappa X^2 + 4\kappa^2 X^2 Y - Y + 2\kappa Y^2) + X(1 - 5\kappa Y - 6\kappa^2 X^2) \tag{3.9}$$

With the ansatz (3.3) the action becomes

$$\mathcal{S}_\mathrm{E} = 2\pi^2 \textstyle\int_0^\infty \mathrm{d}r \; r^3 \; \mathcal{L}_\mathrm{E}(r, \kappa) \tag{3.10a}$$

$$\mathcal{L}_\mathrm{E}(r,\kappa) = 4\kappa^{-2} \left(1 - (1 + \kappa Y)(1 - 2\kappa Y - 3\kappa^2 X^2)^{\frac{1}{2}}\right) \tag{3.10b}$$

One may verify that the equation of motion for this one-dimensional action is identical to eq.(3.9).

If we denote the k dependence of the solution explicitly as $f(r,\kappa)$ then from eq.(3.9) we get the following scale property

$$f(r,\kappa) = f(r/\sqrt{\kappa}, 1) \tag{3.11}$$

In fact by letting $r/\sqrt{\kappa} \to r$ it is easy to see that κ can be removed from the action so that effectively we may take $\kappa = 1$.

We have not been able to obtain an analytic solution of eq.(3.9) for $\kappa \neq 0$. However, it is easy to see that there are three stationary solutions[5]

$$f(r) = \tfrac{1}{2}, \qquad f(r) = 0, \qquad f(r) = 1 \tag{3.12}$$

The action (3.10a) is infinity for $f = \frac{1}{2}$ and zero for $f = 0, 1$.

We define an instanton-like solution as one which interpolates between $f(0) = 0$ and $f(r) \to 1$ as $r \to \infty$. Eq.(3.9) is invariant under $f(r) \to 1 - f(r)$. This relates the instanton-like solution to the anti-instanton-like solution, $f_{\mathrm{a}}(r) = 1 - f_{\mathrm{i}}(r)$.

The same situation occurs for the Yang-Mills action, where the equation of motion is given by eq.(3.9) with $\kappa = 0$

$$0 = f''r^2 + f'r - 4(2f-1)(f-1)f \tag{3.13}$$

The instanton solution for the Yang-Mills action is of the exact form

$$f_{YM}(r) = \frac{r^2}{r^2 + a} \tag{3.14}$$

which may be obtained by solving $X = -Y$.

Now if we write f for large r as

$$f(r) = \lambda + \textstyle\sum_{n=0}^{\infty} a_n r^{\alpha - n} \qquad \alpha < 0 \tag{3.15}$$

and substitute this expression into eq.(3.9) then we get $\alpha = -2$, $\lambda = 0, 1$. Choosing $\lambda = 1$ and $\kappa = 1$ we get an instanton-like solution

$$f_{\mathrm{i}}(r) = 1 - a\frac{1}{r^2} + a^2\frac{1}{r^4} - (a^3 + a^2)\frac{1}{r^6} + \cdots \tag{3.16}$$

where a is a free parameter.

A similar analysis can be done for small r and gives

$$f^{\mathrm{i}}(r) = br^2 - \tfrac{3+20b}{3+12b}b^2r^4 + \tfrac{9+153b+618b^2-464b^3-3584b^4}{9(1-2b)(1+4b)^3}b^3r^6 + \cdots \tag{3.17}$$

where b is a free parameter.

If eqs.(3.16, 3.17) are well defined for $r \geq s$, $r \leq s$ respectively and there exist constants a, b such that $f^{\mathrm{i}}(s) = f_{\mathrm{i}}(s)$, $f^{\mathrm{i}\prime}(s) = f_{\mathrm{i}}'(s)$, then the instanton/anti-instanton-like solutions

---

[5] See the appendix for some analysis of the linearized version of eq.(3.9) around the stationary solutions.

are well defined globally.[6]
With initial conditions, $f'(0)=0,\ f(0)=0$, some numerical analysis[7] support this instanton-like solution:



Fig. 1. INSTANTON-LIKE

As $r \to \infty$, we have $f \to 1$ for the instanton-like solution (3.16) and hence

$$A_\mu \to g^{-1}\partial_\mu g, \qquad\qquad F_{\mu\nu} \to 0 \tag{3.18}$$

In general this condition is sufficient for the integral of the Chern-Pontryagin density, ${}^*F^a_{\mu\nu}F^a_{\mu\nu}$, to be identical with the topological Cartan-Maurer form, since

$${}^*F^a_{\mu\nu}F^a_{\mu\nu} = \partial_\mu \mathrm{tr}(\tfrac{4}{3}\epsilon_{\mu\nu\lambda\rho}A_\nu A_\lambda A_\rho - 4A_\nu{}^*F_{\mu\nu}) \tag{3.19}$$

and the first term gives the Cartan-Maurer form as a surface integral which depends on only the value of f at infinity while the second term vanishes [11]. Thus our instanton-like solution has the same winding number as the ordinary instanton solution in the Yang-Mills action, which is one

$$-\tfrac{1}{32\pi^2}\int \mathrm{d}^4x\ {}^*F^a_{\mu\nu}F^a_{\mu\nu} = 1 \tag{3.20}$$

A similar analysis can be done for the anti-instanton-like solution, where $A_\mu(x)$ has the same singularity at the origin as the ordinary anti-instanton solution of the Yang-Mills theory. Hence the winding number is minus one.

---

[6] In consequence, there will be no free parameter left contrary to the ordinary instanton solutions, where conformal symmetry of the Yang-Mills action allows a free parameter. However, we have not been able to get the exact values of a, b.

[7] We put $\kappa = 1$ for the numerical analysis, Fig. 1, 2, 3.

Some numerical analysis also support this result. Furthermore the action, $\mathcal{S}$, seems[8] to have the exact value, $8\pi^2$.



Fig. 2. Lagrangian



Fig. 3. Action

4 Comment

In four-dimensions the Bianchi identity (3.7) and the equation of motion (2.13) show that ${}^*F^{\mu\nu}$ and $\Pi^{\mu\nu}$ satisfy the same equation. From this observation one may consider solutions of

$$\Pi^{\mu\nu} = c\,{}^*F^{\mu\nu} \tag{4.1}$$

where c is a constant.
Eq.(4.1) reduces to the ordinary self-dual equation for the Yang-Mills action. Hence, one may regard eq.(4.1) as a self-dual equation for general actions.
In the case of SU(2), from the completeness relation (2.3) and eq.(2.12), eq.(4.1) leads to

$$0 = (G+\kappa F)^{-1\mu}{}_{\alpha}{}^{\nu}{}_{\beta} - \frac{1}{2}(G+\kappa F)^{-1\mu}{}_{\gamma}{}^{\nu}{}_{\gamma}\delta_{\alpha\beta} + \kappa c\,({\rm Det}(G+\kappa F))^{-\frac{1}{4}}\,{}^*F^{\mu}{}_{\alpha}{}^{\nu}{}_{\beta} \tag{4.2}$$

Using the identity

$$F^a_{\mu\nu}\,{}^*F^{a\nu\lambda} = -\frac{1}{4}F^a_{\nu\rho}\,{}^*F^{a\nu\rho}\,\delta_\mu{}^\lambda \tag{4.3}$$

we find that in the case of SU(2) eq.(4.1) is equivalent to

$$\left(({\rm Det}(G+\kappa F))^{\frac{1}{4}} - \frac{1}{8}c\kappa^2{\rm tr}(F_{\lambda\rho}{}^*F^{\lambda\rho})\right)F_{\mu\nu} = c\,{}^*F_{\mu\nu} + \frac{1}{2}c\kappa[F_\mu{}^\lambda, {}^*F^\lambda{}_\nu] \tag{4.4}$$

---

[8] We do not have any rigorous proof of it.

However, direct calculation shows that the SO(4) invariant ansatz (3.3) is not compatible with eq.(4.4).

Acknowledgements

The author wishes to thank T. Ahn, H. Osborn, M. Green and A. Mountain at DAMTP, University of Cambridge for valuable discussions. This work was partly supported by Cambridge Overseas Trust and Asia Pacific Center for Theoretical Physics.

Appendix

The linearization of the differential equation (3.9) around the stationary solutions, $f = 0, 1, \frac{1}{2}$ may be achieved by writing $f(r) \simeq f + \delta f(r)$.
Around $f = 0, 1$ we have

$$0 = \delta f'' + \frac{1}{r}\delta f' - \frac{4}{r^2}\delta f \tag{A.5}$$

The solutions are $\delta f = r^{\pm 2}$, which are consistent with eqs.(3.16, 3.17).
Around $f = \frac{1}{2}$ we have

$$0 = (1 - \kappa\frac{1}{2r^2})(1 + \kappa\frac{1}{r^2})\delta f'' + (\frac{1}{r} + \kappa\frac{5}{2r^3})\delta f' + (\frac{2}{r^2} + \kappa\frac{2}{r^4})\delta f \tag{A.6}$$

The asymptotic solution is

$$f(r) \simeq \tfrac{1}{2} + c\sin(\sqrt{2}\ln(r/r_{\text{o}})) \qquad \text{for } r >> 1 \tag{A.7}$$

where c, r_{o} are free parameters.

References

[1] M. Born and L. Infeld. Foundations of the New Field Theory. *Proc. Roy. Soc.*, A144: 425, 1934.

[2] T. Hagiwara. A non-abelian Born-Infeld Lagrangian. *J. Phys.*, A14:3059, 1981.

[3] P. C. Argyres and C. R. Nappi. Spin-1 Effective Actions from Open Strings. *Nucl. Phys.*, B330: 151, 1990.

[4] A. A. Tseytlin . On non-Abelian Generalisation of Born-Infeld Action in String Theory. *Nucl. Phys.*, B501: 41, 1997.

[5] J. Scherk and J. H. Schwarz. Dual Models for Non-Hadrons. *Nucl. Phys.*, B81: 118, 1974.

[6] A. A. Tseytlin. Vector Field Effective Action in the Open Superstring Theory. *Nucl. Phys.*, B276: 391, 1986.

[7] D. Brecher and M. J. Perry. Bound States of D-Branes and the Non-Abelian Born-Infeld Action. hep-th/9801127.

[8] D. Brecher. BPS States of the Non-Abelian Born-Infeld Action. hep-th/9804180.

[9] N. Grandi, E. F. Moreno and F. A. Schaposnik. Monopoles in non-Abelian Dirac-Born-Infeld Theory. hep-th/9901073.

[10] A. A. Belavin, A. M. Polyakov, A. S. Schwartz and Yu. S. Tyupkin. Pseudoparticle Solutions of the Yang-Mills Equations. *Phys. lett.*, 59B: 85, 1975.

[11] S. Weinberg. *The Quantum Theory of Fields.* Cambridge University Press, 1996. See chapter 23 and references therein.